



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

218. Proposed by O. W. ANTHONY, DeWitt Clinton High School, New York City.

From a given triangle cut off an area equivalent to a given square by a line passing through a given point without the triangle.

I. Solution by F. D. POSEY, A. B., Instructor in Mathematics, St. Matthews School, San Mateo, Cal.

Let  $ABC$  be the given triangle and  $P$  the given point on the opposite side  $AC$  say from  $B$ . Let  $a$  be the side of the given square and suppose the part cut off to be a triangle.

Draw  $PE$  perpendicular to  $AC$  and  $PD$  parallel to  $AB$  cutting  $AC$  in  $D$ . Consider the problem solved and the line  $PQR$  cutting  $AC$  and  $AB$  respectively in  $Q$  and  $R$ , to be the required line making  $AQR = a^2$ . Let  $AQ \equiv x$ ,  $QD \equiv y$ ,  $PE \equiv m$ ,  $AD \equiv n$ . Since  $AQR$  and  $PQD$  are similar,

$$\frac{x^2}{y^2} = \frac{AQR}{PQD} = \frac{2a^2}{my}, \quad \frac{x^2}{y^2} = \frac{2a^2}{m}.$$

$$\text{Since } y = n - x, \quad \frac{x^2}{n - x} = \frac{2a^2}{m} \quad \text{or} \quad x^2 + \frac{2a^2}{m}x + \frac{2a^2n}{m}.$$

$$\text{Completing the square, } x^2 + \frac{2a^2}{m}x + \frac{a^4}{m^2} = \frac{a^2}{m} \left( 2n + \frac{a^2}{m} \right).$$

Construct a third proportional  $p$  to  $m$ ,  $a$ .  $\therefore p = a^2/m$ .

$$\therefore x^2 + 2px + p^2 = p[2n + p].$$

Construct  $r = 2n + p$  and  $s$  a mean proportional between  $p$  and  $r$ .

$$\therefore x^2 + 2px + p^2 = s^2, \quad x + p = s, \quad x = s - p.$$

By constructing  $AQ = s - p$ ,  $AQR$  may be shown to have the required area. If the part cut off is not a triangle it may be reduced to the preceding case by cutting off a triangle equivalent to the difference of the given triangle and the given square.

II. Solution by G. W. GREENWOOD, A. M. (Oxon), Professor of Mathematics and Astronomy in McKendree College, Lebanon, Ill.

Lines cutting off a triangle of constant area from two intersecting lines envelope an hyperbola having the given lines as asymptotes.\*

Construct hyperbolas for each pair of sides, and draw tangent to them through the given point.

If we are restricted to compass and ruler, we can construct the foci and the auxiliary circle, and by this means construct the tangents through the points.

Other solutions, which arrived too late for insertion, will be printed next month.

---

\*Take the two lines as the axes of abscissae, the angle between them being  $w$  and the given area  $A$ . The area of the triangle formed by the axes and the line  $x/a + y/b = 1$  is  $\frac{1}{2}absinw = A$ , whence the equation of the line becomes  $(ya^2/2A)sinw - a + x = 0$ . Differentiating with respect to  $a$  and eliminating, the equation of the envelop is found to be the hyperbola  $xy = (1/2sinw)(2A^2 - A)$ . Ed.

III. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let  $ABC$  be the given triangle,  $AB$  base,  $C$  the vertex. Let  $P$  be the given point without,  $PA=b$ ,  $\angle PAC=\theta$ ,  $\angle BAC=A$ , area of square= $m^2$ , area of triangle= $\Delta$ ,  $\Delta - m^2 = n$ . Let the line from  $P$  cutting off the area  $m^2$  meet  $AC$  in  $D$ ,  $AB$  in  $E$ ;  $AD=y$ ,  $AE=x$ .

$$\text{Then } \frac{1}{2}xy\sin A = m^2 \dots\dots\dots (1).$$

$$\frac{1}{2}by\sin\theta = \text{area } PAD \dots\dots\dots (2).$$

$$\frac{1}{2}bx\sin[\theta + A] = \text{area } PAE \dots\dots\dots (3).$$

$$\therefore \frac{1}{2}bx\sin[\theta + A] - \frac{1}{2}by\sin\theta = m^2 \dots\dots\dots (4).$$

The value of  $y$  from (1) in (4) gives

$$x^2 - \frac{2m^2x}{b\sin[\theta + A]} = \frac{2m^2\sin\theta}{\sin A\sin[\theta + A]}.$$

$$\therefore x = \frac{m^2}{b\sin[\theta + A]} \pm \sqrt{\frac{2m^2\sin\theta}{\sin A\sin[\theta + A]} + \frac{m^4}{b^2\sin^2[\theta + A]}},$$

$$y = \frac{\sin[\theta + A]}{\sin\theta} \left[ \pm \sqrt{\frac{2m^2\sin\theta}{\sin A\sin[\theta + A]} + \frac{m^4}{b^2\sin^2[\theta + A]}} - \frac{m^2}{b\sin[\theta + A]} \right].$$

If  $m^2$  is to be the quadrilateral part of the triangle, substitute  $n$  for  $m^2$  in the above values for  $x$ ,  $y$ .

Two geometrical constructions which arrived too late for insertion will be printed next month.

219. On account of the pedagogical importance of this problem, the solutions are withheld until next month in order that they may all be printed in the same number.

# CALCULUS.

Problem 173 was also solved by L. E. Newcomb, Los Gatos, California.

174. Proposed by B. F. FINKEL, A. M., 204 St. Marks Square, Philadelphia, Pa.

Integrate  $\int_0^\infty \frac{\sinh px}{\sinh qx} \cos rx \, dx$ , if  $p^2 < q^2$ .

I. Solution by WILLIAM HOOVER, Ph. D., Professor of Mathematics, Ohio State University, Athens, O.

$$I = \int_0^\infty \frac{\sinh px}{\sinh qx} \cos rx \, dx = \int_0^\infty \frac{e^{px} - e^{-px}}{e^{qx} - e^{-qx}} \cos rx \, dx \dots\dots\dots (1).$$

In (1), put  $qx = \pi z$ , or  $x = [\pi/q]z$ ; then (1) is

$$I = \frac{\pi}{q} \int_0^\infty \frac{e^{az} - e^{-az}}{e^{\pi z} - e^{-\pi z}} \cos m\pi z \, dz \dots\dots\dots (2),$$

in which  $a = p\pi/q$ ,  $m = r\pi/q$ , and the required integral